



# The Physics Potential of the PHENIX VTX and FVTX Detectors

Eric J. Mannel

WWND

13-Apr-2012





# The Relativistic Heavy Ion Collider

- 2 Counter-circulating rings, 3.8 Km circumference
- 200 GeV/Nucleon Au+Au, Cu+Cu, Cu+Au and U+U
- 200/500 GeV polarized p+p
- Search for and study of Quark-Gluon plasma
- Study proton spin








#### PHENIX In RUN-12

- Hadrons, photons and electrons in central arms
  - Drift and Pad chambers for charged particle tracking.
  - Ring Imaging Cerenkov and electromagnetic calorimeter for electron ID
- Muons and Hadrons in the forward regions
  - Mu ID
  - Mu Trackers
  - RPCs RUN 11/12 Upgrade
- VTX for central tracking
- FVTX for forward tracking







#### Heavy Flavor Measurement in PHENIX

- Heavy flavor decays measured indirectly in semileptonic decays
- Measure the spectrum of all electrons
- Subtract background sources
  - Photonic using cocktail of known sources
    - Conversion photons from light meson decays
    - Direct photons from hard scattering process
    - Dalitz decays, p<sup>0</sup>, h, ...
  - Quarkonia contributions
- Separation of B and C decays not possible





## Forward Physics at PHENIX

- Heavy Ion Physics
  - $-J/\Psi$  as a probe of QGP
  - Elliptic Flow
  - Reaction plane: R<sub>AA</sub>
  - Cold Nuclear Effects
- Spin Physics
  - Drell-Yan: Sea quark contribution to proton spin
  - W Program
- Large backgrounds make these measurements difficult





### PHENIX Upgrades

- Recent upgrades to PHENIX tracking
  - Central Silicon Vertex Detector (VTX)
    - Installed in 2010 for RUN-11
  - Forward Silicon Vertex Detector (FVTX)
    - Installed in 2011 for RUN-12
  - Improved vertex resolution
    - Designed to separate heavy decays (charm and bottom)
    - Improved measurements for both Heavy Ion and Spin Physics at PHENIX





# Physics of the VTX

#### • Heavy Ions:

- Heavy flavor production along with beauty and charm separation constrain energy loss models
- Charm measurements along with improved vector meson measurements helps with the understanding of vector meson production and suppression
- Charm and beauty energy flow

#### • Spin:


 Heavy flavor measurements allow understanding of gluon spin contribution.






# PH\*ENIX R<sub>AA</sub> and V2 Measurements


#### Current: PHENIX PRL 98:172301 (2007)







# VTX Gluon Polarization Measurement



- Gluon-Compton process, Photon+ Jet is a "golden" channel
- Large acceptance of VTX allows reconstruction of x from the recoil jet axis
- Allows extending the xrange over current detector.

4/13/2012 E.J. Mannel, BNL 9






## **FVTX Physics**


- Single Muons:
  - Precision heavy flavor and hadron measurements at forward rapidity
  - Separation of charm and beauty
  - W background rejection improved
- Dimuons:
  - Direct measurement of bottom via  $B \rightarrow J/\psi$
  - Separation of  $J/\psi$  from  $\psi'$  with improved resolution.
  - Direct measurement of c-cbar events via  $\mu^+\mu^-$  becomes possible
- Physics:
  - Precise heavy flavor measurements of R<sub>AA</sub> and flow.
  - Detection of  $\psi'$  plus heavy quark allows detailed understanding of vector meson production and modification
  - Precise gluon polarization and sea quark measurements over large x range,





# Expected Improvement in J/Ψ










# Spin Measurement Improvements

- FVTX allows precision measurements for heavy quarks
- Extends the X range in both directions.
- Curves show various estimates of the gluon polarization (T. Gehrmann and W. J. Stirling, Z. Phys. C65, 461 (1995)








## The VTX

#### • 4 Barrels

- Inner 2 barrels, pixels
- Outer 2 barrels, 2-DStrips
- Covers  $|\eta|$  < 1.2
- DCA resolution of 50μm
- Secondary Vertex finding
- b/c separation
- D->Kπ, B->J/ $\Psi$ +X, J/ $\Psi$ ->e<sup>+</sup>e<sup>-</sup>








#### **VTX Current Status**

- Installed prior to RUN-11
- Commissioned during 500 GeV p-p running
- 4.8B Minimum Bias 200 GeV Au-Au events recorded with VTX
- Physics analysis in progress
- RUN-12 in progress:
  - 200 GeV and 500 GeV p-p
  - U-U and Cu-Au runs to come






#### VTX In RUN-11



The reconstructed beam spot using VTX tracks Measured size is 105µm and consistent with expected size

Figures courtesy of R. Nouicer


Typical 200 GeV Au-Au reconstructed event from RUN-11








#### VTX In Run 11



Energy deposition in the strip layers for tracks with  $|\eta| < 0.5$ 

Single track DCA to the beam spot. Estimated DCA resolution is  $88\mu m$  for tracks with a pT > 1GeV




Figures courtesy of R. Nouicer





# VTX In RUN-12




Reconstructed 200 GeV p-p event from RUN-12





#### The FVTX

- 4 planes per end-cap
- Covers
  - $-1.2 < |\eta| < 2.4$
  - $-2\pi$  in  $\varphi$
  - -18.5 cm < |z| < 38 cm
- Resolutuion:
  - Hit < 25 $\mu$ m
  - DCA < 200  $\mu m$







#### **FVTX Status**

- Construction completed and detector installed in 2011
- Commissioning during RUN-12 200/500 GeV p+p running
- Data taking has started in RUN-12:
  - 200/500 GeV p+p
  - 200 GeV U+U
  - 200 GeV Cu+Au
- Data analysis is in progress





#### **FVTX Status**

- Single track projection to the Z-axis
  - From RUN-12 500 GeV pp data
  - Comparison with BBC vertex




Figure courtesy A. Key





# **FVTX Tracking**

- Preliminary hit resolution
- Analysis is still in a very preliminary state
- Will improve as alignment is finalized




Figure courtesy A. Key





#### Conclusions

- Both the VTX and FVTX have been successfully installed
- VTX has taken data for RUN-11 and RUN-12:
- FVTX has taken data for RUN-12
- Analysis of VTX/FVTX data is in progress
- Look for first results from VTX at QM-12, FVTX results will follow quickly after that





## Special Thanks To

Member of the PHENIX VTX and FVTX groups
The PHENIX technical staff
Members of the PHENIX Collaboration
BNL C-AD Department
The Organizers of WWND 2012